

Self sustained phantom wormholes in semi-classical gravity

Remo Garattini*

*Università degli Studi di Bergamo, Facoltà di Ingegneria,
Viale Marconi 5, 24044 Dalmine (Bergamo) ITALY and
INFN - sezione di Milano, Via Celoria 16, Milan, Italy.*

Francisco S. N. Lobo†

*Institute of Gravitation & Cosmology, University of Portsmouth, Portsmouth PO1 2EG, UK and
Centro de Astronomia e Astrofísica da Universidade de Lisboa,
Campo Grande, Ed. C8 1749-016 Lisboa, Portugal*

A possible candidate for the late time accelerated expanding Universe is phantom energy, which possesses rather bizarre properties, such as the prediction of a Big Rip singularity and the violation of the null energy condition. The latter is a fundamental ingredient of traversable wormholes, and it has been shown that phantom energy may indeed sustain these exotic geometries. Inspired by the evolving dark energy parameter crossing the phantom divide, we consider in this work a varying equation of state parameter dependent on the radial coordinate, i.e., $\omega(r) = p(r)/\rho(r)$. We shall impose that phantom energy is concentrated in the neighborhood of the throat, to ensure the flaring out condition, and several models are analyzed. We shall also consider the possibility that these phantom wormholes be sustained by their own quantum fluctuations. The energy density of the graviton one loop contribution to a classical energy in a phantom wormhole background and the finite one loop energy density are considered as a self-consistent source for these wormhole geometries. The latter semi-classical approach prohibits solutions with a constant equation of state parameter, which further motivates the imposition of a radial dependent parameter, $\omega(r)$, and only permits solutions with a steep positive slope proportional to the radial derivative of the equation of state parameter, evaluated at the throat. The size of the wormhole throat as a function of the relevant parameters is also explored.

PACS numbers: 04.62.+v, 04.90.+e

I. INTRODUCTION

Unravelling the source for the late time accelerated expansion of the Universe has become a central theme in modern cosmology [1]. Several candidates, responsible for this expansion, have been proposed in the literature, ranging from modified gravity theories to the introduction of cosmological models comprising of a negative pressure cosmic fluid, denoted as dark energy. The latter is parameterized by the equation of state $\omega = p/\rho$, where p is the spatially homogeneous pressure and ρ the dark energy density. A value of $\omega < -1/3$ is required to ensure the cosmic accelerated expansion, and $\omega = -1$ corresponds to the presence of a cosmological constant. It is interesting that recent constraints from observational data have confirmed that the dark energy parameter lies slightly below the characteristic cosmological constant value, $\omega = -1$ [2]. Ameliorated fits to the data also indicate that an evolving equation of state parameter which crosses $\omega = -1$ is mildly favored [3]. The models satisfying $\omega < -1$ are denoted as *phantom energy* models, and possess rather interesting characteristics, in particular, the prediction of the *Big Rip* singularity and the violation of the null energy condition (NEC). The latter is a fundamental ingredient of traversable wormholes, and it was recently shown that phantom energy may indeed support these exotic geometries [4, 5]. Inspired by the evolving dark energy parameter crossing the phantom divide, we consider in this work a varying equation of state parameter dependent on the radial coordinate, i.e., $\omega(r) = p(r)/\rho(r)$, and impose that phantom energy is concentrated in the neighborhood of the throat, in order to ensure the NEC violation, $\omega(r_0) < -1$, where r_0 is the wormhole throat radius. The latter restriction may be relaxed far from the throat. Despite the fact that the dark (phantom) energy equation of state represents a spatially homogeneous cosmic fluid, due to gravitational instabilities inhomogeneities may arise. Note that now one regards that the pressure in the equation of state $p = \omega\rho$ is a negative radial pressure, and the tangential pressure p_t is determined through the Einstein field equations. This is fundamentally the analysis carried out in Ref. [6], where the authors constructed a time-dependent solution describing a spherically symmetric wormhole

*Electronic address: Remo.Garattini@unibg.it

†Electronic address: floba@cosmo.fis.fc.ul.pt

in a cosmological setting with a ghost scalar field. It was shown that the radial pressure is negative everywhere and far from the wormhole throat equals the transverse pressure, showing that the ghost scalar field behaves essentially as dark energy. Another interesting aspect of this topic is the effect of phantom energy accretion in wormholes immersed in this cosmic fluid. In fact, González-Díaz analyzed the evolution of wormhole and ringhole spacetimes embedded in a background accelerating Universe [7] driven by dark energy, and further considered the accretion of dark and phantom energy onto Morris-Thorne wormholes [8]. It was shown that this accretion gradually increases the wormhole throat which eventually overtakes the accelerated expansion of the universe and becomes infinite at a time in the future before the big rip.

Thus, the phantom wormhole models considered in this work may possibly arise from density fluctuations in the cosmological background [9]. In this context, we shall also be interested in the analysis that these phantom wormholes be sustained by their own quantum fluctuations. As noted above, wormholes violate the NEC, and consequently violate all of the other classical energy conditions. Thus, it seems that these exotic spacetimes arise naturally in the quantum regime, as a large number of quantum systems have been shown to violate the energy conditions, such as the Casimir effect. Indeed, various wormhole solutions in semi-classical gravity have been considered in the literature. For instance, semi-classical wormholes were found in the framework of the Frolov-Zelnikov approximation for $\langle T_{\mu\nu} \rangle$ [10]. Analytical approximations of the stress-energy tensor of quantized fields in static and spherically symmetric wormhole spacetimes were also explored in Refs. [11]. However, the first self-consistent wormhole solution coupled to a quantum scalar field was obtained in Ref. [12]. The ground state of a massive scalar field with a non-conformal coupling on a short-throat flat-space wormhole background was computed in Ref. [13], by using a zeta renormalization approach. The latter wormhole model, which was further used in the context of the Casimir effect [14], was constructed by excising spherical regions from two identical copies of Minkowski spacetime, and finally surgically grafting the boundaries (A more realistic geometry was considered in Ref. [15]). In this context, this motivates the semi-classical analysis in this work, where the Einstein field equation takes the form

$$G_{\mu\nu} = \kappa \langle T_{\mu\nu} \rangle^{\text{ren}}, \quad (1)$$

with $\kappa = 8\pi G$, and $\langle T_{\mu\nu} \rangle^{\text{ren}}$ is the renormalized expectation value of the stress-energy tensor operator of the quantized field. The formalism outlined in this work follows Ref. [16] very closely. Now, the metric may be separated into a background component, $\bar{g}_{\mu\nu}$ and a perturbation $h_{\mu\nu}$, i.e., $g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$. The Einstein tensor may also be separated into a part describing the curvature due to the background geometry and that due to the perturbation, i.e.,

$$G_{\mu\nu}(g_{\alpha\beta}) = G_{\mu\nu}(\bar{g}_{\alpha\beta}) + \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta}), \quad (2)$$

where $\Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}, h_{\alpha\beta})$ may be considered a perturbation series in terms of $h_{\mu\nu}$. Using the semi-classical Einstein field equation, in the absence of matter fields, one may define an effective stress-energy tensor for the quantum fluctuations as

$$\langle T_{\mu\nu} \rangle^{\text{ren}} = -\frac{1}{\kappa} \langle \Delta G_{\mu\nu}(\bar{g}_{\alpha\beta}) \rangle^{\text{ren}}, \quad (3)$$

so that the equation governing quantum fluctuations behaves as a backreaction equation. The semi-classical procedure followed in this work relies heavily on the formalism outlined in Ref. [16], where the graviton one loop contribution to a classical energy in a traversable wormhole background was computed, through a variational approach with Gaussian trial wave functionals [16, 17]. A zeta function regularization is used to deal with the divergences, and a renormalization procedure is introduced, where the finite one loop is considered as a self-consistent source for traversable wormholes. Rather than reproduce the formalism, we shall refer the reader to Ref. [16] for details, when necessary. In this paper, rather than integrate over the whole space as in Ref.[16], we shall work with the energy densities, which provides a more general working hypothesis.

This paper is outlined in the following manner: In Section II, we outline the field equations governing phantom wormholes with an r -dependent parameter, $\omega(r)$, and explore specific solutions. In Section III, we analyze self-sustained traversable phantom wormholes in semi-classical gravity, considering that the equation governing quantum fluctuations behaves as a backreaction equation. Finally, in Section IV, we conclude.

II. FIELD EQUATIONS AND SOLUTIONS

The spacetime metric representing a spherically symmetric and static wormhole is given by

$$ds^2 = -e^{2\Phi(r)} dt^2 + \frac{dr^2}{1 - b(r)/r} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (4)$$

where $\Phi(r)$ and $b(r)$ are arbitrary functions of the radial coordinate, r , denoted as the redshift function, and the form function, respectively [19]. The radial coordinate has a range that increases from a minimum value at r_0 , corresponding to the wormhole throat, to infinity. A fundamental property of a wormhole is that a flaring out condition of the throat, given by $(b - b'r)/b^2 > 0$, is imposed [19, 20], and at the throat $b(r_0) = r = r_0$, the condition $b'(r_0) < 1$ is imposed to have wormhole solutions. Another condition that needs to be satisfied is $1 - b(r)/r > 0$. For the wormhole to be traversable, one must demand that there are no horizons present, which are identified as the surfaces with $e^{2\Phi} \rightarrow 0$, so that $\Phi(r)$ must be finite everywhere.

Using the Einstein field equation, $G_{\mu\nu} = \kappa T_{\mu\nu}$, with $\kappa = 8\pi G$ and $c = 1$, we obtain the following relationships

$$b' = \kappa r^2 \rho, \quad (5)$$

$$\Phi' = \frac{b + \kappa r^3 p_r}{2r^2(1 - b/r)}, \quad (6)$$

$$p'_r = \frac{2}{r} (p_t - p_r) - (\rho + p_r) \Phi', \quad (7)$$

where the prime denotes a derivative with respect to the radial coordinate, r . $\rho(r)$ is the energy density, $p_r(r)$ is the radial pressure, and $p_t(r)$ is the lateral pressure measured in the orthogonal direction to the radial direction.

Now consider the equation of state representing phantom energy, but with a varying parameter $\omega(r)$, given by $p_r(r) = \omega(r)\rho(r)$ with $\omega < -1$ in the phantom regime [4, 5]. The motivation for this analysis arises from a time-dependent crossing of the phantom divide, however, here we shall consider an r -dependent equation of state, with the imposition of phantom energy distribution in the throat neighborhood, with $\omega(r_0) < -1$, to ensure the NEC violation, and consequently the flaring-out condition. This condition may be relaxed far from the throat. Taking into account Eqs. (5)-(6), we have the following condition

$$\Phi'(r) = \frac{b(r) + \omega(r)rb'(r)}{2r^2 [1 - b(r)/r]}. \quad (8)$$

We now have four equations, namely the field equations, i.e., Eqs. (5)-(7), and Eq. (8), with six unknown functions of r , i.e., $\rho(r)$, $p_r(r)$, $p_t(r)$, $b(r)$, $\Phi(r)$ and $\omega(r)$. One may adopt different strategies to construct solutions with the properties and characteristics of wormholes. One may consider a plausible matter-field distribution, by considering convenient stress-energy tensor components, and through the field equations and Eq. (8) determine the metric fields and the function $\omega(r)$.

Alternatively, by appropriately choosing the form function and $\omega(r)$, one may integrate Eq. (8), to determine the redshift function, $\Phi(r)$. As an example, consider the following functions

$$b(r) = r_0 + \gamma^2 r_0 \left(1 - \frac{r_0}{r}\right), \quad (9)$$

$$\omega(r) = -\alpha^2 + \beta^2 \left(1 - \frac{r_0}{r}\right), \quad (10)$$

with $0 < \gamma^2 < 1$, so that one obtains a wormhole solution with $b'(r) = \gamma^2 r_0^2 / r^2$, and at the throat we have $b(r_0) = r_0$ and $b'(r_0) < 1$. Relatively to the function $\omega(r)$, we impose $\omega(r_0) = -\alpha^2 < -1$, and $-1 < \omega(r) = -\alpha^2 + \beta^2 < -1/3$ as $r \rightarrow \infty$. Now, substituting Eqs. (9)-(10) into Eq. (8), we obtain the solution

$$\begin{aligned} \Phi(r) = & -\frac{1 - \gamma^2 \alpha^2}{2(\gamma^2 - 1)} \ln(r - r_0) - \frac{1}{2} \left(1 + \alpha^2 + \frac{\beta^2}{\gamma^2}\right) \ln(r) \\ & + \frac{\beta^2 - \alpha^2 - \frac{\beta^2}{\gamma^2} + \gamma^2}{2(\gamma^2 - 1)} \ln(r - \gamma^2 r_0) + \frac{\beta^2 r_0}{2r} + C. \end{aligned} \quad (11)$$

This corresponds to a non-traversable wormhole solution, due to the presence of an event horizon, as may be readily verified from the first term in the right hand side. Therefore, we fine-tune the parameter $\alpha^2 = 1/\gamma^2$, so that the redshift function finally takes the form

$$\Phi(r) = \frac{1 + \beta^2 + \gamma^2}{2\gamma^2} \ln \left(1 - \frac{\gamma^2 r_0}{r}\right) + \frac{\beta^2 r_0}{2r}, \quad (12)$$

which is now finite throughout the range of r , and where we have considered $C = 0$, without a significant loss of generality. Note that this solution corresponds to an asymptotically flat traversable wormhole, as $b(r)/r \rightarrow 0$ and $\Phi(r) \rightarrow 0$ for $r \rightarrow \infty$.

One may also consider restricted choices for $\Phi(r)$ and $\omega(r)$, and through Eq. (8) determine the form function through the following relationship

$$b(r) = e^{-\Gamma(r_0, r)} \left[r_0 + \int_{r_0}^r \frac{2\Phi'(\bar{r})\bar{r}}{\omega(\bar{r})\bar{r}} e^{\Gamma(r_0, \bar{r})} d\bar{r} \right], \quad (13)$$

with

$$\Gamma(r_0, r) = \int_{r_0}^r \frac{2\Phi'(\bar{r})\bar{r} + 1}{\omega(\bar{r})\bar{r}} d\bar{r}. \quad (14)$$

Now, interestingly enough results may be inferred by imposing a constant redshift function, $\Phi' = 0$, i.e., $b(r) + \omega(r)rb'(r) = 0$, which may be integrated to yield the following form function

$$b(r) = r_0 \exp \left[- \int_{r_0}^r \frac{d\bar{r}}{\omega(\bar{r})\bar{r}} \right]. \quad (15)$$

The latter may also be deduced from Eq. (13), by setting $\Phi'(r) = 0$. Note that for a constant equation of state parameter, ω , the form function is given by $b(r) = r_0(r/r_0)^{-1/\omega}$, which is a solution analyzed in detail in Ref. [5].

For instance, consider the following function

$$\omega(r) = \frac{\alpha^2 \beta^2 (r/r_0)}{\alpha^2 (1 - r/r_0) - \beta^2}, \quad (16)$$

which takes the values $\omega(r_0) = -\alpha^2 < -1$ at the throat, and $\omega(r) = -\beta^2$ as $r \rightarrow \infty$. Now, substituting this specific function in Eq. (15), we deduce the following form function

$$b(r) = r_0 \left(\frac{r}{r_0} \right)^{1/\beta^2} \exp \left[- \left(1 - \frac{r_0}{r} \right) \left(\frac{1}{\beta^2} - \frac{1}{\alpha^2} \right) \right]. \quad (17)$$

Note that if we wish to impose the crossing of the phantom divide the condition $\beta^2 < 1$ needs to be satisfied. However, from the expression of the form function, we verify that in order to have an asymptotically flat spacetime, i.e., $b/r \rightarrow 0$, as $r \rightarrow \infty$, the condition $\beta^2 > 1$ is imposed. Therefore for this particular choice of $\omega(r)$, the solution remains in the phantom regime for all values of r , so that the wormhole is constituted totally of phantom energy. The derivative of the form function evaluated at the throat, is given by $b'(r_0) = 1/\alpha^2 < 1$, consequently satisfying the flaring-out condition.

Now, consider the function given by Eq. (10), then integrating Eq. (15), we deduce the following form function

$$b(r) = r_0 \left[\frac{\beta^2}{\alpha^2} + \frac{r_0}{r} \left(1 - \frac{\beta^2}{\alpha^2} \right) \right]^{1/(\alpha^2 - \beta^2)}. \quad (18)$$

Note that for this case to be an asymptotically flat spacetime, then we need to impose $\alpha^2 - \beta^2 > 1$. This latter restriction imposes that ω remains in the phantom regime, $\omega(r) < -1$ for all r . The derivative of the form function, evaluated at the throat, is given by $b'(r_0) = 1/\alpha^2 < 1$, which also satisfies the flaring out condition. We note that it is possible, in principle, to construct a wide variety of traversable wormholes supported by phantom energy, with an r -dependent equation of state parameter.

III. SELF-SUSTAINED PHANTOM WORMHOLES

In this Section we shall consider the formalism outlined in detail in Ref. [16], where the graviton one loop contribution to a classical energy in a wormhole background is used. The latter contribution is evaluated through a variational approach with Gaussian trial wave functionals, and the divergences are treated with a zeta function regularization. Using a renormalization procedure, the finite one loop energy was considered a self-consistent source for a traversable wormhole. We refer the reader to Ref. [16] for details.

The classical energy is given by

$$H_{\Sigma}^{(0)} = \int_{\Sigma} d^3x \mathcal{H}^{(0)} = -\frac{1}{16\pi G} \int_{\Sigma} d^3x \sqrt{g} R,$$

where the background field super-hamiltonian, $\mathcal{H}^{(0)}$, is integrated on a constant time hypersurface. R is the curvature scalar, and using metric (4), is given by

$$R = -2 \left(1 - \frac{b}{r} \right) \left[\Phi'' + (\Phi')^2 - \frac{b'}{r(r-b)} - \frac{b'r + 3b - 4r}{2r(r-b)} \Phi' \right].$$

We shall henceforth consider a constant redshift function, $\Phi'(r) = 0$, which provides interestingly enough results, so that the curvature scalar reduces to $R = 2b'/r^2$. Thus, the classical energy reduces to

$$H_{\Sigma}^{(0)} = \frac{1}{2G} \int_{r_0}^{\infty} \frac{dr r^2}{\sqrt{1-b(r)/r}} \frac{b(r)}{\omega(r)r^3}, \quad (19)$$

where the condition $b'(r) = -b(r)/[\omega(r)r]$ has been used to eliminate the $b'(r)$ dependence.

We shall also take into account the total regularized one loop energy given by

$$E^{TT} = 2 \int_{r_0}^{\infty} dr \frac{r^2}{\sqrt{1-b(r)/r}} [\rho_1(\varepsilon) + \rho_2(\varepsilon)], \quad (20)$$

where once again, we refer the reader to Ref. [16] for details. The energy densities, $\rho_i(\varepsilon)$ (with $i = 1, 2$), are defined as

$$\rho_i(\varepsilon) = \frac{1}{4\pi} \mu^{2\varepsilon} \int_{\sqrt{U_i(r)}}^{\infty} d\tilde{E}_i \frac{\tilde{E}_i^2}{\left[\tilde{E}_i^2 - U_i(r) \right]^{\varepsilon-1/2}} = -\frac{U_i^2(r)}{64\pi^2} \left[\frac{1}{\varepsilon} + \ln \left(\frac{\mu^2}{U_i} \right) + 2 \ln 2 - \frac{1}{2} \right]. \quad (21)$$

The zeta function regularization method has been used to determine the energy densities, ρ_i . It is interesting to note that this method is identical to the subtraction procedure of the Casimir energy computation, where the zero point energy in different backgrounds with the same asymptotic properties is involved. In this context, the additional mass parameter μ has been introduced to restore the correct dimension for the regularized quantities. Note that this arbitrary mass scale appears in any regularization scheme.

Note that in analyzing the Einstein field equations, one should consider the whole system of equations, which in the static spherically symmetric case also includes the rr -component. The joint analysis of the equations is necessary to guarantee the compatibility of the system. However, in the semi-classical framework considered in this work there is no dynamical equation for the pressure. Nevertheless, one may argue that the semi-classical part of the pressure is known through the equation of state that determines the relation between the energy density and the pressure.

Since a self sustained wormhole must satisfy

$$H_{\Sigma}^{(0)} = -E^{TT}, \quad (22)$$

which is an integral relation, this has to be true also for the integrand, namely the energy density. Therefore, we set

$$\frac{1}{2G} \frac{b(r)}{\omega(r)r^3} = -2 [\rho_1(\varepsilon) + \rho_2(\varepsilon)]. \quad (23)$$

At first sight one can think that the *phantom* region is straightforward to analyze. However, within the approach of Ref. [16], this is not trivial as the main difference is in the integration. In this paper, we shall consider the approach of working with the energy density, while in Ref. [16], the integration over the whole space was considered. In Appendix A, we show that the classical term integrated over all space forbids the appearance of *phantom energy* [18], at least for constant ω .

For this purpose, the Lichnerowicz equations provide the potentials, which are given by

$$\begin{aligned} U_1(r) &= \frac{6}{r^2} \left[1 - \frac{b(r)}{r} \right] - \frac{3}{2r^2} \left[b'(r) - \frac{b(r)}{r} \right] \\ &= \frac{6}{r^2} \left[1 - \frac{b(r)}{r} \right] + \frac{3b(r)}{2r^3} \left[\frac{1}{\omega(r)} + 1 \right], \end{aligned} \quad (24)$$

$$\begin{aligned} U_2(r) &= \frac{6}{r^2} \left[1 - \frac{b(r)}{r} \right] - \frac{3}{2r^2} \left[\frac{-b'(r)}{2} + \frac{b(r)}{r} \right] \\ &= \frac{6}{r^2} \left[1 - \frac{b(r)}{r} \right] + \frac{3b(r)}{2r^3} \left[\frac{1}{3\omega(r)} - 1 \right], \end{aligned} \quad (25)$$

where the relationship $b'(r) = -b(r)/[\omega(r)r]$ has been used once again in the last terms, to eliminate the $b'(r)$ dependence. We refer the reader to Ref. [16] for the deduction of these expressions.

Thus, taking into account Eq. (21), then Eq. (23) yields the following relationship

$$\frac{1}{2G} \frac{b(r)}{\omega(r)r^3} = \frac{1}{32\pi^2\varepsilon} [U_1^2(r) + U_2^2(r)] + \frac{1}{32\pi^2} \left[U_1^2 \ln \left(\left| \frac{4\mu^2}{U_1\sqrt{e}} \right| \right) + U_2^2 \ln \left(\left| \frac{4\mu^2}{U_2\sqrt{e}} \right| \right) \right]. \quad (26)$$

It is essential to renormalize the divergent energy by absorbing the singularity in the classical quantity, by redefining the bare classical constant G as

$$\frac{1}{G} \rightarrow \frac{1}{G_0} + \frac{2}{\varepsilon} \frac{[U_1^2(r) + U_2^2(r)] \omega(r)r^3}{32\pi^2 b(r)}. \quad (27)$$

Using this, Eq. (26) takes the form

$$\frac{1}{2G_0} \frac{b(r)}{\omega(r)r^3} = \frac{1}{32\pi^2} \left[U_1^2 \ln \left(\left| \frac{4\mu^2}{U_1\sqrt{e}} \right| \right) + U_2^2 \ln \left(\left| \frac{4\mu^2}{U_2\sqrt{e}} \right| \right) \right]. \quad (28)$$

Note that this quantity depends on an arbitrary mass scale. Thus, using the renormalization group equation to eliminate this dependence, we impose that

$$\mu \frac{d}{d\mu} \left[\frac{1}{2G_0} \frac{b(r)}{\omega(r)r^3} \right] = \frac{\mu}{32\pi^2} \frac{d}{d\mu} \left[U_1^2 \ln \left(\left| \frac{4\mu^2}{U_1\sqrt{e}} \right| \right) + U_2^2 \ln \left(\left| \frac{4\mu^2}{U_2\sqrt{e}} \right| \right) \right], \quad (29)$$

which reduces to

$$\frac{b(r)}{\omega(r)r^3} \mu \frac{\partial G_0^{-1}(\mu)}{\partial \mu} = \frac{1}{8\pi^2} [U_1^2(r) + U_2^2(r)]. \quad (30)$$

The renormalized constant G_0 is treated as a running constant, in the sense that it varies provided that the scale μ is varying, so that one may consider the following definition

$$\frac{1}{G_0(\mu)} = \frac{1}{G_0(\mu_0)} + \frac{[U_1^2(r) + U_2^2(r)]}{8\pi^2} \frac{\omega(r)r^3}{b(r)} \ln \left(\frac{\mu}{\mu_0} \right). \quad (31)$$

Thus, Eq. (26) finally provides us with

$$\frac{1}{G_0(\mu_0)} = \frac{\omega(r)r^3}{16\pi^2 b(r)} \left[U_1^2(r) \ln \left(\left| \frac{4\mu_0^2}{U_1(r)\sqrt{e}} \right| \right) + U_2^2(r) \ln \left(\left| \frac{4\mu_0^2}{U_2(r)\sqrt{e}} \right| \right) \right]. \quad (32)$$

Now, the procedure that we shall follow is to find the extremum of the right hand side of Eq. (32) with respect to r , and finally evaluate at the throat r_0 , in order to have only one solution (see discussion in Ref. [16]).

To this effect, we shall use the derivative of the potentials, Eqs. (24)-(25), which we write down for self-completeness and self-consistency, and are given by

$$U_1'(r) = -\frac{12}{r^3} \left(1 - \frac{b}{r} \right) + \frac{6(b-b'r)}{r^4} + \frac{3}{2} \left[\frac{b'r-3b}{r^4} \left(\frac{1}{\omega} + 1 \right) - \frac{b}{r^3} \frac{\omega'}{\omega^2} \right], \quad (33)$$

$$U_2'(r) = -\frac{12}{r^3} \left(1 - \frac{b}{r} \right) + \frac{6(b-b'r)}{r^4} + \frac{3}{2} \left[\frac{b'r-3b}{r^4} \left(\frac{1}{3\omega} - 1 \right) - \frac{b}{r^3} \frac{\omega'}{3\omega^2} \right], \quad (34)$$

and once evaluated at the throat take the following form

$$U_1'(r_0) = \frac{3\mathcal{A}}{r_0^3}, \quad \text{with} \quad \mathcal{A} = 2 \left(\frac{1+\omega_0}{\omega_0} \right) - \frac{1}{2\omega_0^2} [(1+3\omega_0)(1+\omega_0) + \omega_0' r_0], \quad (35)$$

$$U_2'(r_0) = \frac{3\mathcal{B}}{r_0^3}, \quad \text{with} \quad \mathcal{B} = 2 \left(\frac{1+\omega_0}{\omega_0} \right) - \frac{1}{2\omega_0^2} [(1+3\omega_0)(1-3\omega_0) + \frac{\omega_0' r_0}{3}], \quad (36)$$

where the relationship $b' = -b/(\omega r)$ has been used to eliminate the b' dependence. We have defined $\omega(r_0) = \omega_0$, and the term $\omega_0' r_0$ is a number as $\omega_0' \propto 1/r_0$. The potentials, Eqs. (24)-(25) evaluated at the throat reduce to

$$U_1(r_0) = \frac{3X}{2r_0^2} \quad \text{with} \quad X = \left(\frac{1+\omega_0}{\omega_0} \right), \quad (37)$$

$$U_2(r_0) = -\frac{3Y}{2r_0^2} \quad \text{with} \quad Y = \left(\frac{3\omega_0-1}{3\omega_0} \right). \quad (38)$$

To determine the extremum of Eq. (32) with respect to r , we shall consider, for convenience, the following definitions

$$\mathcal{F}(r) = \frac{\omega(r)r^3}{b(r)}, \quad \mathcal{F}'(r) = \frac{r^2}{b(r)}[1 + 3\omega(r) + \omega'(r)r], \quad (39)$$

where in the latter expression, $b' = -b/(\omega r)$ is used. Thus, the extremum of Eq. (32) with respect to r , takes the form

$$\mathcal{F}'(r) \left[U_1^2 \ln \left(\frac{|U_1|\sqrt{e}}{4\mu_0^2} \right) + U_2^2 \ln \left(\frac{|U_2|\sqrt{e}}{4\mu_0^2} \right) \right] + \mathcal{F}(r) \left[2U_1U_1' \ln \left(\frac{|U_1|e}{4\mu_0^2} \right) + 2U_2U_2' \ln \left(\frac{|U_2|e}{4\mu_0^2} \right) \right] = 0, \quad (40)$$

where the following relationship

$$\left[U_i^2 \ln \left(\frac{|U_i|\sqrt{e}}{4\mu_0^2} \right) \right]' = 2U_iU_i' \ln \left(\frac{|U_i|e}{4\mu_0^2} \right), \quad (41)$$

has been used.

Now, evaluated at the throat, r_0 , in order to have only one solution, using Eqs. (35)-(36), and the expressions for the potentials, Eqs. (37)-(38), we verify that Eq. (40) provides

$$\frac{1}{4}(1 + 3\omega_0 + \omega_0' r_0) \left[X^2 \ln \left(\frac{3\sqrt{e}X}{8\mu_0^2 r_0^2} \right) + Y^2 \ln \left(\frac{3\sqrt{e}Y}{8\mu_0^2 r_0^2} \right) \right] + \omega_0 \left[X\mathcal{A} \ln \left(\frac{3eX}{8\mu_0^2 r_0^2} \right) - Y\mathcal{B} \ln \left(\frac{3eY}{8\mu_0^2 r_0^2} \right) \right] = 0.$$

Going through the whole tedious, but fairly straightforward, calculation of the throat, we arrive at

$$\bar{r}_0^2 = \frac{3\sqrt{e}}{8\mu_0^2} \Omega, \quad \text{with} \quad \Omega = X^{\frac{A+C}{E}} Y^{\frac{B+D}{E}} \exp \left(\frac{A+B}{2E} \right). \quad (42)$$

where the constants are defined as

$$A = \omega_0 X\mathcal{A}, \quad B = -\omega_0 Y\mathcal{B}, \quad (43)$$

$$C = \frac{1}{4}(1 + 3\omega_0 + \omega_0' r_0)X^2, \quad (44)$$

$$D = \frac{1}{4}(1 + 3\omega_0 + \omega_0' r_0)Y^2, \quad (45)$$

and $E = A + B + C + D$. We may now consider $\omega_0 < -1$ (phantom regime) and $\omega_0' r_0$ as free parameters, and consequently plot r_0 , which is depicted in Fig. 1. We have defined the dimensionless quantity $R_0 = \bar{r}_0 / \left(\sqrt{\frac{3\sqrt{e}}{8\mu_0^2}} \right) = \sqrt{\Omega}$, which is depicted by the surface in Fig. 1. Note that the throat size decreases for increasing values of the parameter $x = \omega_0' r_0$.




FIG. 1: We have defined the dimensionless quantity $R_0 = \bar{r}_0 / \left(\sqrt{\frac{3\sqrt{e}}{8\mu_0^2}} \right) = \sqrt{\Omega}$, which is depicted by the surface. Note that the throat size decreases for increasing values of the parameter $x = \omega_0' r_0$.

Now, restrictions on the parameter range of ω_0 and $\omega_0' r_0$ may also be imposed. The quantity $G_0(\mu_0)$ must be positive, so that Eq. (32) evaluated at the throat, reduces to

$$\frac{1}{G_0(\mu_0)} = \frac{3|\omega_0|\mu_0^2}{8\pi^2\sqrt{e}\Omega} \ln \left[\left(\frac{X}{\Omega} \right)^{X^2} \left(\frac{Y}{\Omega} \right)^{Y^2} \right]. \quad (46)$$

The log must be positive, which places restrictions on the range of $\omega'_0 r_0$. We shall depict $F = \ln[(X/\Omega)^{X^2}(Y/\Omega)^{Y^2}]$ in Fig. 2. Note that only solutions with a steep positive slope proportional to ω'_0 , at the throat, are permitted. As $\omega_0 \rightarrow -1$, we verify that $\omega'_0 r_0$ increases significantly. Solutions with a constant parameter ω , i.e., $\omega'_0 = 0$, are prohibited, which is in agreement with the calculations in Appendix A.




FIG. 2: The log in the definition of $G_0(\mu_0)$ must be positive, which places restrictions on the range of $x = \omega'_0 r_0$. The surface in the figure is given by $F(x, \omega_0) = \ln[(X/\Omega)^{X^2}(Y/\Omega)^{Y^2}]$. Only solutions with a steep positive slope of the parameter $\omega'_0 r_0$, at the throat, are permitted. As ω_0 tends to -1 , we verify that $\omega'_0 r_0$ increases significantly. Note that solutions with a constant parameter ω , i.e., $\omega'_0 = 0$, are prohibited.

An interesting generic constraint on μ/μ_0 may also be deduced. Substituting Eqs. (42) and (46) into Eq. (31), evaluated at the throat, provides the following relationship

$$\frac{1}{G_0(\mu)} = \frac{3|\omega_0|\mu_0^2}{8\pi^2\sqrt{e}\Omega} \ln \left[\left(\frac{X}{\Omega} \right)^{X^2} \left(\frac{Y}{\Omega} \right)^{Y^2} / \left(\frac{\mu}{\mu_0} \right)^{2(X^2+Y^2)} \right]. \quad (47)$$

Now $G_0(\mu)$ must be positive, so that the above relationship places a constraint on μ/μ_0 , namely

$$\mu < \mu_0 \left[\left(\frac{X}{\Omega} \right)^{X^2} \left(\frac{Y}{\Omega} \right)^{Y^2} \right]^{\frac{1}{2(X^2+Y^2)}}. \quad (48)$$

The above results have been based on a variational calculation, concerning the one-loop graviton contribution in a wormhole background. On general grounds, in order for the one-loop contribution to stabilize the wormhole it can only do so at the Planck scale, so that the wormhole throat is expected to be of the Planck order. We have found an upper bound on the mass parameter μ , and we emphasize that we are not interested in computing an exact result of the latter, but in the possibility that phantom wormholes be, in principle, sustained by their own quantum fluctuations. Indeed, we have demonstrated this fact in the context of an r -dependent equation of state parameter within the one-loop graviton contribution approach.

IV. CONCLUSION

Inspired by the evolving dark energy parameter crossing the phantom divide, we have considered in this work a varying equation of state parameter dependent on the radial coordinate, i.e., $\omega(r) = p(r)/\rho(r)$, and imposed that phantom energy is concentrated in the neighborhood of the throat. The pressure in the equation of state is a negative radial pressure, and the tangential pressure p_t is determined through the Einstein field equations. We have also considered the possibility that these phantom wormholes be sustained by their own quantum fluctuations. The energy density of the graviton one loop contribution to a classical energy in a phantom wormhole background, and the finite one loop energy was considered as a self-consistent source for these phantom wormholes. The size of the wormhole throat as a function of the parameters ω_0 and $\omega'_0 r_0$ was further explored, and in order for the one-loop contribution to stabilize the wormhole it can only do so at the Planck scale, so that the wormhole throat is expected to be of the Planck order. In a rather speculative scenario, one may also argue that the accretion of phantom energy by the wormhole gradually increases the wormhole throat to macroscopic size, much in the spirit of Refs. [7, 8].

At first sight one can think that the *phantom* region is straightforward to analyze. However, within the approach of Ref. [16], this is not trivial. Indeed, the main difference is in the integration. In this paper, we have worked with the

energy density, while in Ref. [16], the integration over the whole space was considered. In Appendix A, we show that the classical term integrated over all space forbids the appearance of *phantom* energy for constant ω [18]. However, it is important to emphasize that the semi-classical approach used in this work only rules out self sustained wormholes, with constant ω , concerning the one-loop graviton contribution, and not the whole plethora of phantom wormholes, as the ones analyzed in Refs. [4, 5]. A few words should be spent concerning the difference between the *phantom* and the *dark* energy case. Actually the *dark* energy (not energy density) could be considered, because the classical term strongly forbids *phantom* energy. Nevertheless, the *dark* energy region is completely incompatible with asymptotic flatness [18]. The analysis for $\omega(r)$ is not a trivial task and it depends on a case to case. However, we have shown that interesting solutions do, in fact, exist, namely those that permit solutions with a steep positive slope proportional to the radial derivative of the equation of state parameter, evaluated at the throat. It is rather important to emphasize a shortcoming in the analysis carried in this paper, mainly due to the difficulties of the semi-classical theory of gravity and, in particular to the technical problems encountered. We have considered a semi-classical variational approach using the energy density, which imposes a local analysis to the problem. Thus, we have restricted our attention to the behavior of the metric function $b(r)$, through a variational approach, at the wormhole throat, r_0 . Despite the fact that the behavior is unknown far from the throat, due to the high curvature effects at or near r_0 , the analysis carried out in this work should extend to the immediate neighborhood of the wormhole throat. Therefore, we may conclude that, in principle, phantom wormhole throats with a varying equation of state parameter dependent on the radial coordinate, may be self sustained by their quantum fluctuations within the semi-classical approach outlined in this paper.

Acknowledgements

FSNL was funded by Fundação para a Ciência e Tecnologia (FCT)–Portugal through the research grant SFRH/BPD/26269/2006.

APPENDIX A: COMPUTATION OF THE CLASSICAL TERM

Here, we compute the hamiltonian of Eq. (19) when $\omega(r)$ is a constant. It is easy to show that

$$H_{\Sigma}^{(0)} = \frac{1}{G\omega} \int_{r_0}^{+\infty} dr \left(\frac{r_0}{r} \right)^{\frac{x}{2}} \frac{1}{\sqrt{\left(\frac{r}{r_0} \right)^x - 1}} \quad x := 1 + \frac{1}{\omega}. \quad (\text{A1})$$

In the previous integral there is an extra factor “2” coming from the counting of the universes. We change variable to obtain

$$\frac{2r_0}{G\omega x} \int_0^{+\infty} \frac{dt}{\cosh^{2-2/x}(t)} = \frac{r_0}{G(1+\omega)} B\left(\frac{1}{2}, \frac{1}{1+\omega}\right) = \frac{r_0}{G} A(\omega) \quad (\text{A2})$$

with

$$A(\omega) = \frac{1}{(1+\omega)} B\left(\frac{1}{2}, \frac{1}{1+\omega}\right) = \frac{\sqrt{\pi}}{(1+\omega)} \frac{\Gamma\left(\frac{1}{1+\omega}\right)}{\Gamma\left(\frac{3+\omega}{2+2\omega}\right)}. \quad (\text{A3})$$

In Eq.(A2), we have used the following formula

$$\int_0^{+\infty} dt \frac{\sinh^{\mu} t}{\cosh^{\nu} t} = \frac{1}{2} B\left(\frac{\mu+1}{2}, \frac{\nu-\mu}{2}\right) \quad \begin{cases} \text{Re } \mu > -1 \\ \text{Re } (\mu - \nu) < 0 \end{cases}. \quad (\text{A4})$$

The existence of the integral is submitted to the following condition ($\mu = 0$) $\omega > -1$ and this shows that the phantom region is forbidden.

[1] A. Grant *et al*, “The Farthest known supernova: Support for an accelerating Universe and a glimpse of the epoch of deceleration,” *Astrophys. J.* **560** 49-71 (2001) [arXiv:astro-ph/0104455];

- S. Perlmutter, M. S. Turner and M. White, “Constraining dark energy with SNe Ia and large-scale structure,” *Phys. Rev. Lett.* **83** 670-673 (1999) [arXiv:astro-ph/9901052];
- C. L. Bennett *et al*, “First year *Wilkinson Microwave Anisotropy Probe* (WMAP) observations: Preliminary maps and basic results,” *Astrophys. J. Suppl.* **148** 1 (2003) [arXiv:astro-ph/0302207];
- G. Hinshaw *et al*, “First year *Wilkinson Microwave Anisotropy Probe* (WMAP) observations: The angular power spectrum,” [arXiv:astro-ph/0302217].
- [2] D. N. Spergel *et al*, “Wilkinson Microwave Anisotropy Probe (WMAP) Three Year Results: Implications for Cosmology,” [arXiv:astro-ph/0603449].
- [3] B. Feng, X. Wang and X. Zhang, “Dark energy constraints from the cosmic age and supernovae,” *Phys. Lett.* **B607** 35-41 (2005) [arXiv:astro-ph/0404224];
- A. Upadhye, M. Ishak and P. J. Steinhardt, “Dynamical dark energy: Current constraints and forecasts,” [arXiv:astro-ph/0411803].
- [4] S. Sushkov, “Wormholes supported by a phantom energy,” *Phys. Rev. D* **71**, 043520 (2005) [arXiv:gr-qc/0502084].
- [5] F. S. N. Lobo, “Phantom energy traversable wormholes,” *Phys. Rev. D* **71**, 084011 (2005) [arXiv:gr-qc/0502099];
- F. S. N. Lobo, “Stability of phantom wormholes,” *Phys. Rev. D* **71**, 124022 (2005) [arXiv:gr-qc/0506001].
- [6] S. V. Sushkov and S. W. Kim, “Cosmological evolution of a ghost scalar field,” *Gen. Rel. Grav.* **36**, 1671 (2004) [arXiv:gr-qc/0404037].
- [7] P. F. González-Díaz, “Wormholes and ringholes in a dark-energy universe,” *Phys. Rev. D* **68**, 084016 (2003) [arXiv:astro-ph/0308382].
- [8] P. F. González-Díaz, “Achronal cosmic future,” *Phys. Rev. Lett.* **93** 071301 (2004) [arXiv:astro-ph/0404045];
- P. F. González-Díaz and J. A. Jiménez-Madrid, “Phantom inflation and the ‘Big Trip’,” *Phys. Lett.* **B596** 16-25 (2004) [arXiv:hep-th/0406261];
- P. F. González-Díaz, “On the accretion of phantom energy onto wormholes,” *Phys. Lett.* **B632** 159-161 (2006) [arXiv:astro-ph/0510771];
- P. F. González-Díaz, “Some notes on the Big Trip,” *Phys. Lett.* **B635** 1-6 (2006) [arXiv:hep-th/0607137].
- [9] F. S. N. Lobo, “Van der Waals quintessence stars,” to appear in *Phys. Rev. D*, [arXiv:gr-qc/0610118].
- [10] S. V. Sushkov, “A selfconsistent semiclassical solution with a throat in the theory of gravity,” *Phys. Lett.* **A164**, 33-37 (1992).
- [11] A. A. Popov and S. V. Sushkov, “Vacuum polarization of a scalar field in wormhole space-times,” *Phys. Rev. D* **63**, 044017 (2001) [arXiv:gr-qc/0009028];
- A. A. Popov, “Stress energy of a quantized scalar field in static wormhole space-times,” *Phys. Rev. D* **64**, 104005 (2001) [arXiv:hep-th/0109166].
- [12] D. Hochberg, A. Popov and S. V. Sushkov, “Self-consistent Wormhole Solutions of Semiclassical Gravity,” *Phys. Rev. Lett.* **78**, 2050 (1997) [arXiv:gr-qc/9701064].
- [13] N. R. Khusnutdinov and S. V. Sushkov, “Ground state energy in a wormhole space-time,” *Phys. Rev. D* **65**, 084028 (2002) [arXiv:hep-th/0202068].
- [14] A. R. Khabibullin, N. R. Khusnutdinov and S. V. Sushkov, “Casimir effect in a wormhole spacetime,” *Class. Quant. Grav.* **23** 627-634 (2006) [arXiv:hep-th/0510232].
- [15] N. R. Khusnutdinov, “Semiclassical wormholes,” *Phys. Rev. D* **67**, 124020 (2003) [arXiv:hep-th/0304176].
- [16] R. Garattini, “Self Sustained Traversable Wormholes?” *Class. Quant. Grav.* **22** 1105-1118 (2005) [arXiv:gr-qc/0501105].
- [17] R. Garattini, “Probing foamy spacetime with variational methods,” *Int.J.Mod.Phys. A* **14** 2905-2920 (1999) [arXiv:gr-qc/9805096];
- R. Garattini, “Vacuum Energy, Variational Methods and the Casimir Energy,” *Phys. Rev. D* **59** 104019 (1999) [arXiv:hep-th/9902006].
- [18] R. Garattini, “Self sustained traversable wormholes and the equation of state,” *Class. Quant. Grav.* **24**, 1189 (2007) [arXiv:gr-qc/0701019].
- [19] M. S. Morris and K. S. Thorne, “Wormholes in spacetime and their use for interstellar travel: A tool for teaching General Relativity,” *Am. J. Phys.* **56**, 395 (1988).
- [20] Visser M 1995 *Lorentzian Wormholes: From Einstein to Hawking* (American Institute of Physics, New York).